


TwoBare. U. Dan Boat]



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 212 945 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
12.06.2002 Bulletin 2002/24

(51) Int Cl.7: **A23C 9/152**, A23C 9/154,
A23C 11/10

(21) Application number: 01310221.5

(22) Date of filing: 06.12.2001

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 07.12.2000 US 251778 P
20.11.2001 US 989617

(71) Applicant: McNeil-PPC, Inc.
Skillman, NJ 08558-9418 (US)

(72) Inventors:
• Boyer, Marie
Fort Washington, PA 19034 (US)

- Hardie-Muncy, Darlene
Ringoos, NJ 08551 (US)
- Hirasuna, Thomas Jyun
Ithaca, NY 14850 (US)
- Myers, Scott
Lansing, NY 14882 (US)
- Roden, Allan
Nobelsville, IN 46060 (US)
- Sharma, Shri Kant
Ithaca, NY 14850 (US)

(74) Representative: Mercer, Christopher Paul et al
Carpmaels & Ransford
43, Bloomsbury Square
London WC1A 2RA (GB)

(54) **Dairy-based beverages fortified with cholesterol lowering agents**

(57) A dairy-based beverage including a stanol ester in an amount of from about 0.2 to about 2.0 wt. % of the beverage and, optionally, an additional ingredient selected from a stabilizer, a weighing agent, an emulsi-

fier, and mixtures thereof, wherein each additional ingredient is in an amount of from about 0.01 to about 0.05 wt % of the beverage.

EP 1 212 945 A2

Description**FIELD OF THE INVENTION**

5 [0001] This invention relates to the incorporation of cholesterol-lowering agents into dairy-based beverages.

Background of the Invention

10 [0002] Human clinical studies have shown that stanol esters in the diet significantly lower cholesterol levels in individuals, thus further reducing the risk of coronary heart disease. On September 8, 2000, the U.S. Food and Drug Administration began authorizing the use, on food labels and in food labeling, of health claims on the association between plant sterol/stanol esters and reduced risk of coronary heart disease (CHD). 65 FR 54685, September 8, 2000. In particular, it was determined that ingestion of 3.4 grams of stanol ester every day for at least two weeks was proven to lower LDL cholesterol levels up to 14% without an adverse effect on HDL levels.

15 [0003] The properties of phytosterols and especially those of stanol esters make them particularly unsuited toward formulation into traditional beverage forms. The poor wettability of these compounds makes them difficult to incorporate into aqueous based forms such as liquids. Additionally, their high melting point and waxy properties make them difficult to process.

20 [0004] Stanol ester is more readily dispersed into fat containing products, as 42% of the standard ingredient is considered fat. Thus, incorporating stanol ester into dairy beverages with lower fat levels, especially reduced fat/fat free milks, is problematic. In addition, the dietary ingredient must be handled in an appropriate manner for its introduction into milk. Once the dietary ingredient has been introduced into milk, formulation and process must be controlled in such a manner as to keep the dietary ingredient in suspension throughout a 60 day shelf life.

25 [0005] Therefore, there is an unmet consumer need for a palatable, stable dosage form for these ingredients in dairy beverages.

SUMMARY OF THE INVENTION

30 [0006] We have found that stanol esters may be successfully incorporated into dairy beverages. This approach results in a palatable, stable dosage form for cholesterol lowering agents. The present invention is a dairy-based beverage including a stanol ester in an amount of from about 0.2 to about 2.0 wt. % of the beverage and, optionally, an additional ingredient selected from a stabilizer, a weighing agent, an emulsifier, and mixtures thereof, wherein each additional ingredient is in an amount of from about 0.01 to about 0.05 wt % of the beverage.

DETAILED DESCRIPTION

35 [0007] The present invention provides an extended shelf-life product with acceptable palatability and mouthfeel characteristics, wherein the cholesterol lowering agent is dispersed throughout the milk matrix. The milk matrix contains among, other things, milk and a cholesterol lowering agent. Optionally, other ingredients are incorporated into the milk matrix, such as, but not limited to emulsifiers, stabilizers, fat, colors, flavors, flavor masking agents, sweeteners, weighing agents and the like.

40 [0008] According to the present invention, "milk" includes whole milk, skim milk, 2% milk, and 1 % milk, as well as soy milk.

45 [0009] While this invention encompasses stanol ester, sterol ester is also within the scope of this invention.

[0010] In addition to milk, other dairy beverages, including, but not limited to, milk shakes, frappes, smoothies, malteds, dairy-based prepackaged liquor beverages, dairy-based prepackaged coffee beverages, complete nutritional dairy beverages, dairy-based diet beverages, and dairy-based infant formulas.

50 [0011] A variety of techniques known to those having skill in the art can be used to disperse stanol ester into milk. However, process capabilities on hand will determine the best dispersion technique. For example, cold or warm stanol ester can be added directly to a cold milk batch with a high shear, in-tank homogenizer; stanol ester can be injected into the milk in front of an inline high shear mixer and recirculated with the milk; in the case of pre-pasteurization, melted stanol ester can be injected into the warm flow in front of the homogenizer during the start of the pasteurization process; shredded stanol ester can be added to about 140°F milk and mixed with a high shear mixer, and mixed or recirculated with the milk in a tank, and melted stanol ester can be injected into preheated milk directly in front of the homogenizer for the UHT process. The stanol ester can be dispersed and hydrated directly into cold (about 40°F) milk with using high shear equipment, e.g., Liquifier agitation.

55 [0012] Stabilizers suitable for use in the present invention are those capable of providing a prolonged dispersion of stanol ester in the milk matrix, preferably for at least 60 days for UHT and longer if aseptically packed. Preferably, the

stabilizer is an edible hydrophilic colloid, i.e., a starch (including pre-gelatinized starch and chemically-modified starch), a gum (either natural or synthetic), or a chemically-modified polysaccharide. Preferred stabilizers suitable for use in the present invention include, but are not limited to, acacia, agar, algin, carrageenan, e.g., CM-611 (FMC Corp., Philadelphia, PA), gum arabic, gum ghatti, gum karaya, gum tragacanth, guar gum, e.g., TIC Guarcel 302 (TIC Gums Inc., Belcamp, MD), locust bean gum, e.g., Meyprodyn 200L (Rhodia Foods, Cranbury, NJ), xanthan gum, sodium alginate, sodium steryl lactolate, xanthan gum, cellulosic gums, such as, methyl cellulose, hydroxy ethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, and carboxymethylcellulose, and mixtures thereof, with carrageenan being preferred. In addition, emulsifier salts, e.g., Joha KM2 (B.K. Landenburg Corp., Simi Valley, CA), weighting agents, e.g., SAIB FG (Eastman Chemical Co., Kingsport, TN), emulsifiers, e.g., K Emplex, SSL (American Ingredient Co., Kansas City, MO), Alphadim 90 NLK (American Ingredient Co., Kansas City, MO), crystal inhibitors, e.g., Caprol ET (AC Humco, Memphis, TN) and mixtures thereof may optionally be added.

[0013] Additionally, it may be possible to eliminate the need for stabilizers by substituting stanol with other selected fatty acids for a stabilizing ingredient.

[0014] According to the present invention, stanol ester is present in the dairy beverage from about 0.2 to about 2.0 wt %, preferably from about 0.2 to about 0.7 wt %. Additional ingredients that may be included in the formulation include from about 0.01 to about 0.05 wt % of a stabilizer, weighing agent, and emulsifier.

[0015] The use of ultra high temperature ("UHT") sterilization is not necessary to carry out the present invention. Although the combination of UHT treatment and refrigerated (38 to 45°F) storage enables a high quality, long-shelf life milk beverage, excellent quality would be expected from same formula processed with typical industry methods of pasteurization (HTST or LTLT techniques) for shorter shelf life products.

[0016] After dispersion, UHT sterilization can be carried out by any known method to those having skill in the art. Either direct or indirect UHT processing may be utilized. As mentioned above, while UHT processing provides for a longer shelf life, e.g., at least 60 days, milk beverage, it is not a necessary requirement for a milk beverage where shorter shelf life, e.g., less than 60 days, is acceptable. Where a shorter shelf life milk beverage is desired, standard milk industry pasteurization techniques provide for a higher quality product, e.g., better flavors, less thermal damage, and the like.

[0017] After UHT processing or pasteurization, the dairy beverage product can be near aseptically filled into purepak containers, preferably at from about 40 to about 70°F. After filling, all subsequent holding, warehousing, distribution, and retailing is done under refrigerated conditions (38 to 45°F). Preferably, the dairy beverage of the instant invention contains skim or 2% milk, stanol ester and carrageenan.

[0018] Briefly, a process for making the 'an extended shelf-life product' of the present invention involves sufficient heating/agitation to facilitate 100% dispersion of stanol ester and stabilizer, e.g., carrageenan, UHT sterilization via direct or indirect steam injection and near aseptic filling of purepak, glass or plastic, containers. All subsequent holding, warehousing, distribution and retailing should be under refrigerated conditions (38-45°F). Controlled temperatures will result in overall better quality.

[0019] If a shorter shelf-life product is desired then standard pasteurization would be utilized instead of UHT sterilization. For longer shelf life, UHT and aseptic packaging may be used.

[0020] One particular method of producing product according to the present invention involved melting stanol ester at about 140°F in a vessel. In a separate vessel, stabilizers, e.g., gums and/or salts, were mixed into milk. The milk-stabilizer mixture was then heated to about 145°F. After heating, stanol ester was added to the heated milk-stabilizer mixture. The resulting mixture was then mixed with a Braun handheld mixer for about 2 minutes. Following hand mixing, two-stage homogenization occurred at about 140°F and 2000-2500/500 psi. Pasteurization at 190°F for 2 minutes was carried out prior to hot-filling sanitized containers. The filled containers were then cooled to less than about 50 °F, followed by refrigerated storage (about 38 to about 45°F).

[0021] After the invention has been described in general hereinbefore, the following examples are intended to illustrate details of the invention, without thereby limiting it in any matter.

Examples

Example 1

[0022] 0.71 % stanol ester (1.7 g stanol ester /80z. serving) 2% milk products containing the stabilizers in Table 1 were prepared as follows: stanol ester (Raisio) was melted and dispersed in skim milk (at about 145°F), and mixed for about 2 minutes with a Braun mixer (Model No. 4169). Stabilizers and salts were then added to the solution and mixed with a Braun mixer (Model No. 4169) for 2 minutes. Two stage homogenization was then conducted at either 2000/500 2500/500 psi. When raw milk was the starting component, the milk was pasteurized at 190°F for 2 minutes. The hot milk was filled into sterilized containers. After packing, the containers were then transferred to less than about 20°F ice-cold water. The cooled containers were then stored in the refrigerator at from about 38 to about 45 °F for sensory

and stability data.

[0023] Stability tests on the product were conducted by visual inspection. The carrageenan containing milks were considered to be the most stable of all examined. Stability, viscosity, accelerated stability measurement (Quickscan) and assessment of additional stabilizers are continuing.

[0024] After two weeks of storage at 40°F, stability tests on the product using a QuickSCAN analyzer (Part No. 6605413, Beckman Coulter Inc., Fullerton, CA). Measuring the percentage of back-scattering referenced to a variable at day 0 provided the ability to identify the amount of creaming.

[0025] The scale used to determine creaming is as follows:

none - 0
slight - less than 10% back-scattering
moderate - 11-20% back-scattering
high - more than 20% back-scattering

TABLE 1,

2% Milk	
Stabilizer(s)	Scan Observations
0.03% carrageenan + 0.1% Meyprodyn 200L	Slight
0.03% carrageenan	Slight
0.3% Guarcel 302, Guar gum	High
0.1% Joha KM2, Emulsifier salt	Moderate
0.3% Joha KM2, Emulsifier salt	High
0.01% carrageenan	Slight
0.0115% carrageenan + 0.0344% SAIB FG, weighing agent	Slight
0.0115% carrageenan + 0.114% Joha KM2	Slight
0.015% carrageenan	Slight
0.02% carrageenan	Slight
0.0175% carrageenan	Slight
0.015% carrageenan + 0.01% Joha KM2 Emulsifier Salt	Slight
0.0175% carrageenan + 0.02% Joha KM2 Emulsifier Salt	Slight

TABLE 1, (continued)

2% Milk	
<u>Stabilizer(s)</u>	<u>Scan Observations</u>
0.0175% carrageenan + 0.05% Joha KM2 Emulsifier Salt	Slight

EXAMPLE 2

[0026] The procedure described in Example 1, above, was followed for the stabilizers described in Table 2, except that skim milk was used.

TABLE 2,

Skim Milk	
<u>Stabilizer(s)</u>	<u>Scan Observations</u>
0.02% carrageenan+0.7066% stanol ester	None
0.1% sodium steryl stearate	High
0.1% mono-diglycerides (Alphadin 90 NLK)	Slight
0.1% polyglycerol esters (Caprol ET)	High

EXAMPLE 3

[0027] The procedure described in Example 1, above, was followed for the stabilizers described in Table 3, except that 1% milk was used.

TABLE 3,

1% Milk	
<u>Stabilizer(s)</u>	<u>Scan Observations</u>
0.02% carrageenan	None

Claims

1. A dairy-based beverage comprising stanol ester in an amount of from about 0.2 to about 2.0 wt. % of the beverage.
2. A dairy-based beverage of claim 1 further comprising an additional ingredient selected from the group consisting of a stabilizer, a weighing agent, an emulsifier, and mixtures thereof, wherein each additional ingredient is in an amount of from about 0.01 to about 0.05 wt % of the beverage.
3. A dairy-based beverage of claim 2 wherein the stabilizer is selected from the group consisting of an edible hydrophilic colloid, a gum, a modified polysaccharide, and mixtures thereof.
4. A dairy-based beverage of claim 3 wherein the stabilizer is selected from the group consisting of acacia, agar, algin, carrageenan, gum arabic, gum ghatti, gum karaya, gum tragacanth, guar gum, locust bean gum, xanthan gum, sodium alginate, sodium steryl lactolate, xanthan gum, cellulosic gums, such as methyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, and carboxymethylcellulose, and mixtures thereof.

5. A dairy-based beverage of claim 4 wherein the stabilizer is carrageenan.
6. A dairy-based beverage of claim 1 wherein the dairy-based beverage is selected from the group consisting of milk, milk shakes, frappes, smoothies, malteds, dairy-based prepackaged liquor beverages, dairy-based prepackaged coffee beverages, complete nutritional dairy beverages, dairy-based diet beverages, and dairy-based infant formulas.
7. A milk comprising from about 0.2 to about 0.7 wt% of a stanol ester and 0.01 to about 0.05 wt% of a stabilizer.
8. A milk of claim 7 wherein the stabilizer is carrageenan.
9. A milk of claim 7 wherein the milk is selected from the group consisting of whole milk, skim milk, 2% milk, 1% milk, and soy milk.
10. A milk of claim 7 further comprising from about 0.01 to about 0.05 wt % of a weighing agent, from about 0.01 to about 0.05 wt % an emulsifier, or both.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 212 945 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
20.11.2002 Bulletin 2002/47

(51) Int Cl.7: **A23C 9/152**, A23C 9/154,
A23C 11/10, A23F 5/24,
A23L 1/30

(43) Date of publication A2:
12.06.2002 Bulletin 2002/24

(21) Application number: **01310221.5**

(22) Date of filing: **06.12.2001**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **07.12.2000 US 251778 P**
20.11.2001 US 989617

(71) Applicant: **McNeil-PPC, Inc.**
Skillman, NJ 08558-9418 (US)

(72) Inventors:
• **Boyer, Marie**
Fort Washington, PA 19034 (US)

- **Hardie-Muncy, Darlene**
Ringoes, NJ 08551 (US)
- **Hirasuna, Thomas Jyun**
Ithaca, NY 14850 (US)
- **Myers, Scott**
Lansing, NY 14882 (US)
- **Roden, Allan**
Nobelsville, IN 46060 (US)
- **Sharma, Shri Kant**
Ithaca, NY 14850 (US)

(74) Representative: **Mercer, Christopher Paul et al**
Carpmaels & Ransford
43, Bloomsbury Square
London WC1A 2RA (GB)

(54) **Dairy-based beverages fortified with cholesterol lowering agents**

(57) A dairy-based beverage including a stanol ester in an amount of from about 0.2 to about 2.0 wt. % of the beverage and, optionally, an additional ingredient selected from a stabilizer, a weighing agent, an emulsi-

fier, and mixtures thereof, wherein each additional ingredient is in an amount of from about 0.01 to about 0.05 wt % of the beverage.

EP 1 212 945 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 01 31 0221

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 00 41491 A (VULFSON EVGENY NAUM ; NUTRAHEALTH LTD (GB); LAW BARRY ARNOLD (GB)) 20 July 2000 (2000-07-20) * claims 1,14,29; examples 18,30 *	1,6	A23C9/152 A23C9/154 A23C11/10 A23F5/24 A23L1/30
X	WO 99 15546 A (ANTTILA ERKKI ; HEIKKILAE ELINA (FI); LINDEMAN ANNELI (FI); MIKKONE) 1 April 1999 (1999-04-01) * example 31 *	1,6	
X	WO 00 45648 A (FORBES MEDI TECH INC) 10 August 2000 (2000-08-10) * page 7, paragraph 2; example 5 *	1,6	
X	FR 2 775 620 A (GERVAIS DANONE CO) 10 September 1999 (1999-09-10) * claims 1-5; examples 1-4 *	1,6	
A	US 6 087 353 A (WALLIS SIMON HOWARD ET AL) 11 July 2000 (2000-07-11) * column 15, line 1 - line 3; claims 1,2,9 *	1,6,7	
A	T. NGUYEN: "The cholesterol-lowering action of plant stanol esters" THE JOURNAL OF NUTRITION, vol. 129, no. 12, 1999, pages 2109-2112, XP008007936 * the whole document *	1,7	A23C A23F A23L
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26 September 2002	Examiner Desmedt, G
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technical background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on or after the filing date D : document cited in the application I : document cited for other reasons S : member of the same patent family, corresponding document	

EPQ FORM 1501 (3-02) (P3) (Cv)

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 01 31 0221

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

26-09-2002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0041491	A	20-07-2000	AU	1995400 A	01-08-2000
			EP	1146798 A2	24-10-2001
			WO	0041491 A2	20-07-2000
WO 9915546	A	01-04-1999	FI	973647 A	10-03-1999
			FI	974563 A	29-04-1999
			FI	974648 A	06-05-1999
			AU	9164198 A	12-04-1999
			AU	9164298 A	12-04-1999
			EP	1015476 A1	05-07-2000
			EP	1027368 A1	16-08-2000
			WO	9915546 A1	01-04-1999
			WO	9915547 A1	01-04-1999
			JP	2002516816 T	11-06-2002
			JP	2001517678 T	09-10-2001
			US	6441206 B1	27-08-2002
WO 0045648	A	10-08-2000	AU	2426500 A	25-08-2000
			WO	0045648 A1	10-08-2000
			EP	1148793 A1	31-10-2001
			US	2002048606 A1	25-04-2002
FR 2775620	A	10-09-1999	FR	2775620 A1	10-09-1999
			AT	217155 T	15-05-2002
			AU	2731999 A	20-09-1999
			CA	2320365 A1	10-09-1999
			DE	69901433 D1	13-06-2002
			DK	1059851 T3	26-08-2002
			EP	1059851 A1	20-12-2000
			WO	9944442 A1	10-09-1999
			JP	2002505093 T	19-02-2002
US 6087353	A	11-07-2000	AU	3596599 A	06-12-1999
			WO	9959421 A1	25-11-1999
			EP	1082022 A1	14-03-2001

EPC FORM P-X/59

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)